



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

AVERAGE AND PROBABILITY,

Conducted by B.F.FINKEL, Kidder, Missouri. All contributions to this department should be sent to him.


SOLUTIONS OF PROBLEMS.

20. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

A surface one inch square is thrown at random upon a surface one foot square, but in such a manner as always to lie wholly upon the larger surface. Find the mean value of the sum of the distances of the vertices of the smaller surface, from any vertex of the larger surface.

Solution by the PROPOSER.

Let $S_1 S_2 = s$, $A_1 A_2 = 12s$, $AD = x$, $DO = y$; then $S_1 O = \frac{1}{2}s\sqrt{2}$. Put $\angle S_1 O P_1 = \angle S_2 O P_2 = \angle S_3 O P_3 = \angle S_4 O P_4 = \theta$; then $DD' = \frac{1}{2}s\sqrt{2} \sin \theta$, and $(DO - P_1 D') = \frac{1}{2}s\sqrt{2} \cos \theta$.



$\therefore A_1 D' = \frac{1}{2}(2x - s\sqrt{2} \sin \theta)$, and $P_1 D' = \frac{1}{2}(2y - s\sqrt{2} \cos \theta)$; also,
 $\Delta_1 = A_1 P_1 = \frac{1}{2}\sqrt{[(2x - s\sqrt{2} \sin \theta)^2 + (2y - s\sqrt{2} \cos \theta)^2]}$, $\Delta_2 = A_1 P_2$
 $= \frac{1}{2}\sqrt{[(2x - s\sqrt{2} \cos \theta)^2 + (2y + s\sqrt{2} \sin \theta)^2]}$, $\Delta_3 = A_1 P_3 = \frac{1}{2}\sqrt{[(2x + s\sqrt{2} \sin \theta)^2 + (2y + s\sqrt{2} \cos \theta)^2]}$, $\Delta_4 = A_1 P_4 = \frac{1}{2}\sqrt{[(2x + s\sqrt{2} \cos \theta)^2 + (2y - s\sqrt{2} \sin \theta)^2]}$.

Let $\Delta = (\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4)$; then five cases confront us, for consideration.

First Case—The point O may lie in the square surface $A_1 A_2 A_3 A_4$. If $\frac{1}{2}(24 - \sqrt{2})s = a$ and $\frac{1}{2}s\sqrt{2} = b$, the mean value of the sum of the distances in this case becomes

$$D_1 = \int_b^a \int_b^a \int_0^{2\pi} [\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4] dx dy d\theta \div \int_b^a \int_b^a \int_0^{2\pi} dx dy d\theta \dots (1).$$

Second Case—The point O may lie in the rectangular surface D_2 ; then if $11\frac{1}{2}s = c$, $\frac{1}{2}s = e$, and $\sin^{-1}(e/e_1/2) = \phi$, the mean value in this case becomes

$$D_2 = \int_c^e \int_e^c \int_0^\phi [\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4] dx dy d\theta \div \int_c^e \int_e^c \int_0^\phi dx dy d\theta \dots (2).$$

Third Case—The point O may lie in the rectangular surface D_3 ; and in this case,

$$D_3 = \int_e^b \int_b^c \int_0^\phi [\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4] dx dy d\theta \div \int_e^b \int_b^c \int_0^\phi dx dy d\theta \dots (3).$$

Fourth Case—The point O may lie in the rectangular surface D_4 . Put $\frac{1}{2}s(23 - \sqrt{2}) = f$, and $\frac{1}{2}s(\sqrt{2} + 1) = g$; then the mean value in this case becomes

$$D_4 = \int_g^f \int_f^b \int_0^\phi [\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4] dx dy d\theta \div \int_g^f \int_f^b \int_0^\phi dx dy d\theta \dots (4).$$

Fifth Case—The point O may lie in the rectangular surface D_5 ; then the mean value in this case becomes

$$D_5 = \int_g^f \int_f^c \int_0^\phi [\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4] dx dy d\theta \div \int_g^f \int_f^c \int_0^\phi dx dy d\theta \dots (5).$$

Consequently the *required* mean value becomes

$$D = \frac{1}{5}(D_1 + D_2 + D_3 + D_4 + D_5) \dots (6);$$

and the labor required in the performance of the integrations indicated is simply enormous.

21. Proposed by H. W. DRAUGHON, Clinton, Louisiana.

From one corner of a square field, a boy runs in a random direction, with a random uniform velocity. The greatest distance the boy can run in one minute is equal to the diagonal of the field. What is the probability that the boy will be in the field at the end of one minute?

Solution by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Let $AB = a$, and $BP = x$; then $AP = \sqrt{(x^2 + a^2)} = w$, and $AC = a\sqrt{2} = m$. The boy will be in the field at the expiration of $t = 1$ minute, if v be not greater than $\sqrt{(x^2 + a^2)}$. Hence the required probability becomes

$$P = \int_0^a \int_0^w dx dv \div \int_0^a \int_0^m dx dv = \frac{1}{a^2 \sqrt{2}} \int_0^a \frac{1}{\sqrt{(x^2 + a^2)}} dx$$

